



Access this article online

Quick Response Code:



Website:
www.turkjemerged.com

DOI:
10.4103/tjem.tjem_175_23

Invited commentary on “The impact of the first wave of the COVID-19 pandemic on hospital admissions and treatment management of ectopic pregnancy”

Fikri M. Abu-Zidan*

The Research Office, College of Medicine and Health Sciences, United Arab Emirates University, Al-Ain, United Arab Emirates
*Corresponding author

I had the privilege to review with great interest the paper of Gurbuz *et al.*^[1] which has been published in the current issue of *Turkish Journal of Emergency Medicine*. I congratulate the authors for their effort in trying to answer an important question. The research design is acceptable, and the statistical analysis is adequate. Their retrospective descriptive study of a single center in Turkey compares the admitted women with ectopic pregnancy during the first 10 weeks of the COVID-19 pandemic with the 114 weeks before the pandemic. There were 116 patients during the first period (one patient per week) compared with 57 patients during the 10 early weeks of the pandemic (5.7 patients per week). The delay of the ectopic pregnancy diagnosis resulted in more ruptures during the pandemic compared with the pre-pandemic period (28.1% compared with 11.2%). Laparotomy was applied to all patients during the pandemic while more than 70% had laparoscopy before the pandemic. This follows the early recommendations of the World Society of Emergency Surgery which indicates that laparoscopy is a high-risk aerosol-generating procedure^[2] which may increase the risk of infection. The

retrospective nature of the study, the small sample size, being from a single center, and the finding of a dramatic increase of the cases (by 6 times) without a clear explanation weaken the strength of the findings.

The management of infectious disasters has four phases which are preparedness, mitigation, response, and recovery.^[3,4] The short COVID-19 studied period of the current study^[1] covers the mitigation (reducing the impact by restricting the movement) and the response phases and may have selection bias. It would have been better to have a longer study period with a larger number of patients (multicentric or national data) which would permit a proper time series analysis to study changes over time than just compare two periods.

A recent systematic review and meta-analysis^[5] have shown that delivery during the COVID-19 pandemic slightly increased from 48% to 54% of all obstetrical and gynecological hospital admissions including an increase of ectopic pregnancy from 1.8% to 2%. Patients, being afraid of COVID-19 infection, would avoid seeking health care unless really needed.^[5] Fear was more during the early stages of

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow_reprints@wolterskluwer.com

How to cite this article: Abu-Zidan FM. Invited commentary on “The impact of the first wave of the COVID-19 pandemic on hospital admissions and treatment management of ectopic pregnancy”. *Turk J Emerg Med* 2023;23:244-5.

Submitted: 10-08-2023
Accepted: 12-08-2023
Published: 03-10-2023

ORCID:
FMAZ: 0000-0003-0143-8568

Address for correspondence:

Prof. Fikri M. Abu-Zidan,
The Research Office,
College of Medicine
and Health Sciences,
United Arab Emirates
University, Al-Ain,
United Arab Emirates.
E-mail: fabuzidan@uaeu.
ac.ae



the pandemic because of the severity of the disease, high infection rate, and lack of effective treatment or vaccine.^[6] The lockdown and stay-home policy reduced the mobility of the community. We have recently shown the reduced number of road traffic collisions and trauma hospitalization in our setting during the pandemic. Nevertheless, those hospitalized trauma patients had more severe injuries with increased hospital deaths.^[7,8]

The increased percentage of ectopic pregnancies reported in the current study must be considered with caution. This increase was not related to increased pregnancies. There was a large drop in births associated with conceptions during the first several months of the pandemic.^[9] Although COVID-19 infection is associated with worse pregnancy outcomes,^[10] it is unlikely, by common sense, to have a direct effect on the pathology of ectopic pregnancy itself. Furthermore, the authors did not report the percentage of those who had COVID-19 infection comparing them with non-CVOD-19 patients.

Finally, I would like to thank the authors for addressing an important area. Nevertheless, I disagree that “it is impossible to carry out a prospective investigation on this concept.” I am confident that prospective cohort studies are possible within infectious disease disasters as demonstrated by one of our recent global studies.^[11] Research should be an integral part of preparedness and response plans for any infectious disease disaster which will produce more evidence that can help us plan the mitigation, response, and recovery of future disasters.

Author contribution statement

Fikri Abu-Zidan reviewed the published manuscript, read the literature, and wrote the commentary.

Conflicts of interest

None declared.

Funding

None.

References

- Gurbuz H, Basol G, Altintas MM, Kuru B. The impact of the first wave of the COVID-19 pandemic on hospital admissions and treatment management of ectopic pregnancy. *Turk J Emerg Med* 2023. Doi: 10.4103/tjem.tjem_37_23.
- De Simone B, Chouillard E, Sartelli M, Biffi WL, Di Saverio S, Moore EE, *et al*. The management of surgical patients in the emergency setting during COVID-19 pandemic: The WSES position paper. *World J Emerg Surg* 2021;16:14.
- Khan G, Sheek-Hussein M, Al Suwaidi AR, Idris K, Abu-Zidan FM. Novel coronavirus pandemic: A global health threat. *Turk J Emerg Med* 2020;20:55-62.
- Sheek-Hussein M, Alsuwaidi AR, Davies EA, Abu-Zidan FM. Monkeypox: A current emergency global health threat. *Turk J Emerg Med* 2023;23:5-16.
- Carbone L, Raffone A, Travaglino A, Saccone G, Di Girolamo R, Neola D, *et al*. The impact of COVID-19 pandemic on obstetrics and gynecology hospitalization rate and on reasons for seeking emergency care: A systematic review and meta-analysis. *J Matern Fetal Neonatal Med* 2023;36:2187254.
- Sheek-Hussein M, Abu-Zidan FM, Stip E. Disaster management of the psychological impact of the COVID-19 pandemic. *Int J Emerg Med* 2021;14:19.
- Yasin YJ, Grivna M, Abu-Zidan FM. Global impact of COVID-19 pandemic on road traffic collisions. *World J Emerg Surg* 2021;16:51.
- Yasin YJ, Alao DO, Grivna M, Abu-Zidan FM. Impact of the COVID-19 pandemic on road traffic collision injury patterns and severity in Al-Ain City, United Arab Emirates. *World J Emerg Surg* 2021;16:57.
- National Bureau of Economic Research. How Did the COVID Pandemic Affect the Number of Births? *The Digest*; 2022. Available from: <https://www.nber.org/digest/202207/how-did-covid-pandemic-affect-number-births>. [Last accessed on 2023 Aug 09].
- Jeong Y, Kim MA. The coronavirus disease 2019 infection in pregnancy and adverse pregnancy outcomes: A systematic review and meta-analysis. *Obstet Gynecol Sci* 2023;66:270-89.
- De Simone B, Abu-Zidan FM, Chouillard E, Di Saverio S, Sartelli M, Podda M, *et al*. The ChoCO-W prospective observational global study: Does COVID-19 increase gangrenous cholecystitis? *World J Emerg Surg* 2022;17:61.