



Access this article online

Quick Response Code:



Website:

<https://turkjemergmed.com/>

DOI:

10.4103/tjem.tjem\_73\_24

Submitted: 12-04-2024

Revised: 05-06-2024

Accepted: 05-06-2024

Published: 01-07-2024

ORCID:

AŞ: 0000-0002-0583-2936

MP: 0000-0002-3917-0192

MM: 0000-0001-6928-2307

EA: 0000-0002-3249-2420

SK: 0000-0002-8692-7266

GA: 0000-0002-5580-3201

NÖD: 0000-0002-5209-8076

BÖ: 0000-0002-6645-7287

MH: 0000-0001-6368-6853

ÇT: 0009-0001-5534-5781

SEE: 0000-0002-3183-3713

YY: 0000-0003-3917-9434

NA: 0000-0001-8059-4862

NGG: 0000-0001-5052-9242

NNU: 0000-0003-3509-3254

HA: 0000-0002-8396-4854

ASG: 0000-0003-2669-0532

SK: 0000-0001-8265-9615

SA: 0000-0002-2197-4715

AÖ: 0000-0002-2707-4099

GPG: 0000-0001-8531-4591

MBS: 0000-0001-9335-9001

ÖS: 0000-0002-5557-6627

SBHB: 0000-0002-7695-

9969

**Address for  
correspondence:**

Dr. Alp Şener,  
Department of Emergency  
Medicine, Faculty of  
Medicine, Ankara Bilkent  
City Hospital, Ankara  
Yıldırım Beyazıt University,  
Bilkent, Ankara, Türkiye.  
E-mail: alpsener@  
gmail.com

# Prospective, multicenter, Turkish out-of-hospital cardiac arrest study: TROHCA

Alp Şener<sup>1,2\*</sup>, Murat Pekdemir<sup>3</sup>, Mehmet Muzaffer İslam<sup>4</sup>, Ersin Aksay<sup>5</sup>, Sevilay Karahan<sup>6</sup>, Gokhan Aksel<sup>4</sup>, Nurettin Özgür Doğan<sup>3</sup>, Berkant Öztürk<sup>7</sup>, Muhammet Hacimustafaoğlu<sup>7</sup>, Çağrı Türkücü<sup>7</sup>, Serkan Emre Eroğlu<sup>4</sup>, Yusuf Yürümez<sup>8</sup>, Nuray Aslan<sup>8</sup>, Necip Gokhan Güner<sup>8</sup>, Neşe Nur User<sup>9</sup>, Hüseyin Aldemir<sup>9</sup>, Abdullah Sadık Girişgin<sup>10</sup>, Sedat Koçak<sup>10</sup>, Sami Ataman<sup>10</sup>, Ayhan Özhasenekler<sup>1,2</sup>, Gul Pamukçu Günaydın<sup>1,2</sup>, Mustafa Burak Sayhan<sup>11</sup>, Ömer Salt<sup>11</sup>, Satuk Bugra Han Bozatlı<sup>11</sup>, Engin Deniz Arslan<sup>12</sup>, Fevzi Yılmaz<sup>12</sup>, Ramazan Sivil<sup>12</sup>, Özlem Köksal<sup>13</sup>, Vahide Aslıhan Durak<sup>13</sup>, Fatma Özdemir<sup>13</sup>, Mahmut Taş<sup>14</sup>, Yenal Karakoç<sup>14</sup>, Öner Avınca<sup>14</sup>, Yunus Emre Arık<sup>15</sup>, Adem Melekoğlu<sup>15</sup>, Özgür Çevrim<sup>15</sup>, Özlem Yiğit<sup>16</sup>, Cem Oktay<sup>16</sup>, Süleyman İbze<sup>16</sup>, Salim Satar<sup>17</sup>, Muge Gülen<sup>17</sup>, Selen Acehan<sup>17</sup>, Erhan Altunbaş<sup>18</sup>, Melis Efeoğlu Saçak<sup>18</sup>, Emir Ünal<sup>18</sup>, Erdem Çevik<sup>19</sup>, Dilay Satılmış<sup>19</sup>, Hande Asan<sup>19</sup>, Yunus Karaca<sup>20</sup>, Melih İmamoğlu<sup>20</sup>, Vildan Özer<sup>20</sup>, Ahmet Demircan<sup>21</sup>, Ayfer Keleş<sup>21</sup>, Gültekin Kadı<sup>21</sup>, Orhan Delice<sup>22</sup>, Sibel Güçlü Utlu<sup>22</sup>, Senol Arslan<sup>22</sup>, Neslihan Yücel<sup>23</sup>, Şükrü Gürbüz<sup>23</sup>, Hüseyin Burak Ayhan<sup>23</sup>, Abdullah Şen<sup>24</sup>, Mahmut Yaman<sup>24</sup>, Müge Günalp<sup>25</sup>, Sinan Genç<sup>25</sup>, Ahmet Baydın<sup>26</sup>, Fatih Çalışkan<sup>26</sup>, Şeyma Arzu Temür<sup>26</sup>, Murat Ersel<sup>27</sup>, Sercan Yalçınlı<sup>27</sup>, Enver Özçete<sup>27</sup>, Bulent Erbil<sup>28</sup>, Elif Ozturk Ince<sup>28</sup>, Mehmet Ali Karaca<sup>28</sup>, Murat Çetin<sup>29</sup>, Mehmet Demirbağ<sup>30</sup>, Mustafa Sabak<sup>31</sup>, Mustafa Bozkurt<sup>31</sup>

<sup>1</sup>Department of Emergency Medicine, Ankara Bilkent City Hospital, Ankara, Türkiye, <sup>2</sup>Department of Emergency Medicine, Ankara Yıldırım Beyazıt University, Faculty of Medicine, Ankara, Türkiye, <sup>3</sup>Department of Emergency Medicine, Kocaeli University, Faculty of Medicine, Kocaeli, Türkiye, <sup>4</sup>University of Health Sciences, Umraniye Training and Research Hospital, İstanbul, Türkiye, <sup>5</sup>Department of Emergency Medicine, İzmir Ekonomi University, Medical Point Hospital, İzmir, Türkiye, <sup>6</sup>Department of Biostatistics, Hacettepe University, Faculty of Medicine, Ankara, Türkiye, <sup>7</sup>Department of Emergency Medicine, Aksaray University, Aksaray Training and Research Hospital, Aksaray, Türkiye, <sup>8</sup>Department of Emergency Medicine, Sakarya University, Faculty of Medicine, Sakarya, Türkiye, <sup>9</sup>Department of Emergency Medicine, Afyonkarahisar Health Sciences University, Afyonkarahisar, Türkiye, <sup>10</sup>Department of Emergency Medicine, Necmettin Erbakan University, Faculty of Medicine, Konya, Türkiye, <sup>11</sup>Department of Emergency Medicine, Trakya University, Faculty of Medicine, Edirne, Türkiye, <sup>12</sup>Department of Emergency Medicine, University of Health Sciences, Antalya Training and Research Hospital, Antalya, Türkiye, <sup>13</sup>Department of Emergency Medicine, Bursa Uludağ University, Faculty of Medicine, Bursa, Türkiye, <sup>14</sup>Department of Emergency Medicine, University of Health Sciences, Gazi Yaşargil Training and Research Hospital, Diyarbakır, Türkiye, <sup>15</sup>Department of Emergency Medicine, University of Health Sciences, İstanbul Şişli Hamidiye Etfal Training and Research Hospital, İstanbul, Türkiye, <sup>16</sup>Department of Emergency Medicine, Akdeniz University, Faculty of Medicine, Antalya, Türkiye, <sup>17</sup>Department of Emergency Medicine, University of Health Sciences, Adana City Training and Research Hospital, Adana, Türkiye, <sup>18</sup>Department of Emergency Medicine, Marmara University Pendik Training and Research Hospital, İstanbul, Türkiye, <sup>19</sup>Department of Emergency Medicine, University of Health Sciences, Sultan 2. Abdülhamid Han Training and Research Hospital, İstanbul, Türkiye, <sup>20</sup>Department of Emergency Medicine, Karadeniz Technical University, Faculty of Medicine, Trabzon, Türkiye, <sup>21</sup>Department of Emergency Medicine, Gazi University, Faculty of Medicine, Ankara, Türkiye, <sup>22</sup>Department of Emergency Medicine, Erzurum City Hospital, Erzurum, Türkiye, <sup>23</sup>Department of Emergency Medicine, İnönü University, Turgut Özal Medical Center, Malatya, Türkiye, <sup>24</sup>Department of Emergency Medicine, Dicle University, Faculty of Medicine, Diyarbakır, Türkiye, <sup>25</sup>Department of Emergency Medicine, Ankara University, Faculty of Medicine, Ankara, Türkiye, <sup>26</sup>Department of Emergency Medicine, Ondokuz Mayıs University, Faculty of Medicine, Samsun, Türkiye, <sup>27</sup>Department of Emergency

This is an open access journal, and articles are distributed under the terms of the Creative Commons Attribution-NonCommercial-ShareAlike 4.0 License, which allows others to remix, tweak, and build upon the work non-commercially, as long as appropriate credit is given and the new creations are licensed under the identical terms.

For reprints contact: WKHLRPMedknow\_reprints@wolterskluwer.com

**How to cite this article:** Şener A, Pekdemir M, İslam MM, Aksay E, Karahan S, Aksel G, *et al.* Prospective, multicenter, Turkish out-of-hospital cardiac arrest study: TROHCA. Turk J Emerg Med 2024;24:133-44.

EDA: 0000-0002-0155-6903 *Medicine, Ege University, Faculty of Medicine, İzmir, Türkiye,* <sup>28</sup>*Department of Emergency Medicine, Hacettepe University, Faculty of Medicine, Ankara, Türkiye,* <sup>29</sup>*Dr.Behçet Uz Pediatric Diseases and Surgery Training and Research Hospital, İzmir, Türkiye,* <sup>30</sup>*Department of Emergency Medicine, Manisa Merkezefendi State Hospital, Manisa, Türkiye,* <sup>31</sup>*Department of Emergency Medicine, Gaziantep University, Şahinbey Research and Practice Hospital, Gaziantep, Türkiye*  
 FY: 0000-0002-3675-7457  
 RS: 0000-0002-6463-2117  
 ÖK: 0000-0003-2271-5659  
 VAD: 0000-0003-0836-7862  
 FÖ: 0000-0002-4278-1576  
 MT: 0000-0002-3894-4766  
 YK: 0000-0003-3607-9286  
 ÖA: 0000-0003-3109-6773  
 YEA: 0000-0002-4521-9546  
 AM: 0000-0002-5764-3678  
 ÖÇ: 0000-0002-5515-5232  
 ÖY: 0000-0002-3703-7590  
 CO: 0000-0001-7993-0566  
 Sl: 0000-0002-0927-0029  
 SS: 0000-0001-6080-4287  
 MG: 0000-0002-5080-3501  
 SA: 0000-0003-4390-6593  
 EA: 0000-0002-5075-739X  
 MES: 0000-0002-8446-9898  
 EÜ: 0000-0003-2758-8214  
 EÇ: 0000-0003-1474-3166  
 DS: 0000-0003-3765-2208  
 HA: 0000-0002-6742-302X  
 YK: 0000-0002-0947-0565  
 MI: 0000-0003-4197-8999  
 VÖ: 0000-0002-2514-5674  
 AD: 0000-0002-7993-8098  
 AK: 0000-0003-4429-3100  
 GK: 0000-0002-9195-6635  
 OD: 0000-0003-1629-4245  
 SGU: 0000-0001-8635-1562  
 SA: 0000-0002-6636-5307  
 NY: 0000-0001-5845-2614  
 ŞG: 0000-0003-2616-0304  
 HBA: 0000-0002-8876-1378  
 AŞ: 0000-0001-6705-4079  
 MY: 0000-0002-0783-1419  
 MG: 0000-0001-8140-6720  
 SG: 0000-0002-0516-1028  
 AB: 0000-0003-4987-0878  
 FÇ: 0000-0001-7786-3929  
 ŞAT: 0009-0003-6564-2434  
 ME: 0000-0003-2282-5559  
 SY: 0000-0003-1780-8391  
 EÖ: 0000-0002-1685-2369  
 BE: 0000-0001-8555-0017  
 EÖl: 0000-0002-6396-6668  
 MAK: 0000-0003-0876-2811  
 MÇ: 0000-0002-0667-7966  
 MD: 0000-0002-0475-872X  
 MS: 0000-0003-2777-2003  
 MB: 0000-0002-4400-6696

**Abstract:**

**OBJECTIVES:** There is no sufficient data to provide a clear picture of out-of-hospital cardiac arrest (OHCA) across Türkiye. This study is the first to present the prognostic outcomes of OHCA cases and the factors associated with these outcomes.

**MATERIALS AND METHODS:** The study was conducted in a prospective, observational, multicenter design under the leadership of the Emergency Medicine Association of Turkey Resuscitation Study Group. OHCA cases aged 18 years and over who were admitted to 28 centers from Türkiye were included in the study. Survived event, return of spontaneous circulation (ROSC), survival to hospital discharge, and neurological outcome at discharge were investigated as primary outcomes.

**RESULTS:** One thousand and three patients were included in the final analysis. 61.1% of the patients were male, and the average age was 67.0 ± 15.2. Cardiopulmonary resuscitation (CPR) was performed on 86.5% of the patients in the prehospital period by emergency medical service, and bystander CPR was performed on only 2.9% by nonhealth-care providers. As a result, the survived event rate was found to be 6.9%. The survival rate upon hospital discharge was 4.4%, with 2.7% of patients achieving a good neurological outcome upon discharge. In addition, the overall ROSC and sustained ROSC rates were 45.2% and 33.4%, respectively. In the multiple logistic regression analysis, male gender, initial shockable rhythm, a shorter prehospital duration of CPR, and the lack of CPR requirement in the emergency department were determined to be independent predictors for the survival to hospital discharge.

**CONCLUSION:** Compared to global data, survival to hospital discharge and good neurological outcome rates appear to be lower in our study. We conclude that this result is related to low bystander CPR rates. Although not the focus of this study, inadequate postresuscitative care and intensive care support should also be discussed in this regard. It is obvious that this issue should be carefully addressed through political moves in the health and social fields.

**Keywords:**

Bystander cardiopulmonary resuscitation, cardiopulmonary resuscitation, cardiopulmonary resuscitation, out-of-hospital cardiac arrest, registry, return of spontaneous circulation, survival, survived event, Turkey, Türkiye

**Box-ED section**

**What is already known about the study topic?**

- Out-of-hospital cardiac arrest (OHCA) remains a global public health problem. Although there are varying rates, survival rates are far from satisfactory. Türkiye’s data on this subject are limited; therefore, a multicenter study was needed.

**How is this study structured?**

- The study was planned under the supervision of the Emergency Medicine Association of Turkey Resuscitation Study Group for 1-year period with a prospective, observational, multicenter design in OHCA patients aged 18 years and over.

**What does this study tell us?**

- In this study, the rate of survival to

hospital discharge for OHCA was found to be 4.4%; there are similar results in the literature. However, it is thought provoking that bystander cardiopulmonary resuscitation (CPR) was applied only in 2.9% of the patients.

**What is the conflict on the issue? Is it important for readers?**

- Bystander-initiated CPR is one of the key points of a chain of survival. In our country, where the prehospital system and emergency medicine are well developed, the low rate of hospital discharge of out-of-cardiac arrest victims may be due to the low bystander CPR rate. Therefore, improved policies are needed in the political and health fields to enable the public to early recognize OHCA and start early bystander-initiated CPR.

## Introduction

The overall incidence of out-of-hospital cardiac arrests (OHCAs) where cardiopulmonary resuscitation (CPR) was attempted was 56 per 100,000 population annually.<sup>[1]</sup> Although it is a frequently discussed issue in the literature, it remains an important public health problem. In particular, the American Heart Association (AHA) carries out regular updates of their recommendations.<sup>[2]</sup> Even if these guidelines are followed, survival data in OHCA cases are far from satisfactory globally and vary widely. These variations may be related to the design of the studies and the lack of standardization in the definitions used. To improve the results, it is necessary to expand these data, determine strategic goals with newly emerging data, and adapt them to life with the support of health-care and social policies. Assessing each step of the survival chain individually, particularly early recognition of cardiac arrest, prompt activation of emergency medical services (EMS), and enhancing bystander CPR rates, can be regarded as strategic objectives aimed at bolstering the survival rates.<sup>[3]</sup> It is known that bystander CPR rates are not sufficient worldwide and even this rate is far from expectations in Türkiye.

Data on this subject are limited in Türkiye. Generally, there are retrospective, single-center studies, as well as a few prospective studies with low sample size.<sup>[4-6]</sup> With these limited data, it is not possible to talk about the generalizability of the data such as survival, return of spontaneous circulation (ROSC), and bystander CPR. It is known that monitoring and keeping arrest records influence survival. With this prospective, multicenter study, we aimed to fill this data gap, shed light on future studies, and guide health policies.

## Materials and Methods

### Settings

This prospective, observational, multicenter study was planned to be conducted for 1 year (2023) in 30 health centers in Türkiye within the scope of the Emergency Medicine Association of Turkey Resuscitation Study Group. All study centers were third-level emergency department (ED) providing emergency medicine training, and these centers were selected from all 7 geographical regions of Türkiye, and all resuscitation practices are carried out in accordance with the AHA and/or European Resuscitation Council guidelines. Two centers were excluded from the study because data could not be obtained. Ultimately, the study was completed in 28 centers with a total of more than 8 million ED visits annually [Figure 1].

There is generally no specific protocol used for terminating resuscitation, and the decision depends on the physician's initiative. EMS system of Türkiye generally operates on the "load and go" principle. In the field, EMS staff rarely make end-of-life decisions, and the majority of patients are transported to the emergency department (ED). It has been accepted that this principle was followed throughout this study. We should also note that it is still not a legal obligation in Türkiye to install automated external defibrillators (AEDs), especially in all densely crowded places such as malls and bus stations.

### Population

Patients who had a cardiac arrest in an out-of-hospital environment, were 18 years of age or older, and had ROSC before admission or were admitted as cardiac




Figure 1: Distribution of the study centers

arrest to the ED were included 24/7 in the study. Informed consent was obtained from all patients/relatives. Patients experiencing traumatic cardiac arrest, those transferred to study centers from other hospitals, defined as deceased prior to admission, whose primary outcome data couldn't be obtained, and whose relatives did not provide consent for their participation in the study, were excluded from the study. According to the sample size calculation, based on the ROSC (24.8%) and survival to hospital discharge (5.6%) rates in the study of Şener *et al.* and assuming a 5% error, a total of at least 837 cases were planned to be included in the study.<sup>[6]</sup> Upon reaching this sample, the study was terminated on October 15, 2023, before the planned date.

Ethical approval was received from the Kocaeli University Non-Interventional Clinical Research Ethics Committee (No: KÜ GOKAEK-2022/18.24) at the coordinator center on November 10, 2022, and the study was carried out in accordance with the principles of the World Medical Association Declaration of Helsinki.

### Data collection

Data obtained from the patients themselves or their relatives and recorded forms in Hospital Information Management Systems were checked and transferred to the online registry form (Google® Forms) by the local coordinator of the relevant center. Afterward, the relevant data were processed by the authorized researchers and directed for statistical analysis.

### Definitions and outcomes

The data collection forms were primarily designed based on the Utstein Resuscitation Registry Templates for out-of-hospital cardiac arrest (OHCA), with additional variables incorporated.<sup>[7]</sup> Obtaining any rhythm with pulse for at least 5 min during resuscitation was defined as ROSC, and persistence of this condition for at least 20 min was defined as sustained ROSC. Patients who were admitted to the ED with a pulse that persisted 20 min or who completed 20 min with a pulse in the ED were defined as a survived event. Patients who were discharged with sustained ROSC were recorded as survival to hospital discharge. Patients who were still in ROSC at day 30 were also recorded as 30-day survival. Cerebral performance category (CPC) was evaluated at discharge and on the 30<sup>th</sup> day, and according to this scale, categories 1 and 2 were recorded as good neurological outcome and 3 and above were recorded as poor neurological outcome. The hospital parameter as cardiac arrest localization refers to patients who had cardiac arrest at the hospital entrance (parking lot, garden, entrance of the outpatient clinics or triage, etc.) before being delivered to the relevant health-care professional. A cardiac arrest which is seen by other people or is monitored by EMS is defined as witnessed arrest. CPR procedure initiated by witnessed


people other than the EMS staff is defined as bystander CPR. The rhythm of the patient detected in the first monitoring was recorded as initial cardiac arrest rhythm, and pulseless ventricular tachycardia and ventricular fibrillation rhythms were recorded as shockable rhythm. The concept of airway procedure was used as the most advanced method performed. The concept of unsuccessful airway procedure refers to a patient who cannot be ventilated in any way. Survived event, ROSC, survival to hospital discharge, and neurological outcome at discharge were investigated as primary outcomes.

### Statistical analysis

Statistical analyses of the study were performed with IBM SPSS Statistics for Windows, Version 26.0 (Armonk, NY, USA: IBM Corp). In comparing continuous data between two independent groups, the Mann-Whitney *U*-test or Independent samples *t*-test was used, depending on distribution. Continuous data are expressed as mean with standard deviation or median with interquartile range in accordance with distribution. The comparison of categorical data between independent groups was performed with Pearson's Chi-square and Fisher's exact tests; these data were expressed as sample numbers and percentages. In primary analyses,  $P < 0.05$  level was used for statistical significance. Multiple logistic regression analysis was performed for sustained ROSC and discharge, and variables with  $P < 0.250$  in univariate analysis but thought to be clinically significant were included in the model. For discharge outcome, "prehospital CPR, arrest cause, prehospital airway management, arrest location, use of magnesium sulfate in the emergency department (ED), and experiencing cardiac arrest again in the ED" were unable to be included in the model due to imbalances in sample counts across groups. Similarly, for sustained ROSC, "prehospital CPR, arrest etiology, prehospital airway management, and CPR administered in the ED" couldn't be included due to group imbalances. As a result, odds ratios were expressed with 95% confidence intervals. Cutoff values of pH variable used in logistic regression analysis were determined by ROC analysis.

### Results

A total of 1060 patients were recorded in the data system from 28 centers in Türkiye. After ruling out missing data and samples that did not meet the criteria (such as trauma patients, inhospital arrests, and referrals from other hospitals), 1002 patients were included in the final analysis [Figure 2]. It was determined that 61.1% of the patients were male, the mean age was  $67.0 \pm 15.2$  years, the etiology was acute coronary syndrome in 46.3%, and the localization of cardiac arrest was "home" in 61.3%. Witnessed arrest was detected in 83.9% of patients, transport with EMS in 97.4%, and shockable first arrest rhythm in 10.8%. CPR was performed on 86.5% of the



EMS: Emergency medical services, OHCA: Out-of-hospital cardiac arrest, CPR: Cardio-pulmonary resuscitation, ROSC: Return of spontaneous circulation, ED: Emergency department

Figure 2: Flow diagram

patients in the prehospital period by EMS personnel, and bystander CPR was performed on only 2.9% by nonhealth-care providers (higher in male gender; 3.8 vs. 1.5). While the prehospital bag-valve-mask rate was 31.7%, successful endotracheal intubation (ETI) was found to be 47.6%. Peripheral intravenous access was presented in 88.4% of the patients, and 11.2% were delivered to the ED without any venous access [Table 1].

### Primary outcomes

As a result, while the prehospital ROSC rate was 13.6%, the survived event rate was found to be 6.9% [Table 1].

While the mortality rate in the ED was 72.3%, 4.4% were discharged alive from the hospital and 2.7% were discharged with good neurological outcome. In addition, the ROSC rate for the entire process was determined as 45.2% and the sustained ROSC rate was 33.4%.

### Secondary outcomes

In the EDs, mechanical chest compression device was used in 41.5% of patients, EtCO<sub>2</sub> monitoring in 30.1%, and ultrasound-guided CPR performed in 52.1% of patients [Table 2]. Primary comparative

**Table 1: Demographic characteristics and prehospital descriptive variables**

Variables	n (%) or mean±SD or median (25%–75%)
Time zone	
Day shift	386 (38.5)
Night shift and holidays	616 (61.5)
Gender	
Female	390 (38.9)
Male	612 (61.1)
Age (year), mean±SD	67.0±15.2
Etiology	
Acute coronary syndrome	464 (46.3)
Electrolyte disorder	51 (5.1)
Pulmonary embolism	43 (4.3)
Asthma	16 (1.6)
Drowning	16 (1.6)
Intoxication	9 (0.9)
Stroke	9 (0.9)
Electrocution	3 (0.3)
Anaphylaxis	2 (0.2)
Postprimary coronary intervention	2 (0.2)
Postbypass	2 (0.2)
Others	140 (14)
Unknown	245 (24.5)
Independent living	721 (77.4)
Chronic neurological disease	168 (17.9)
Chronic pulmonary disease	183 (19.4)
Chronic cardiovascular disease	496 (52.3)
Active malignancy	126 (13.4)
Ventricular assist device	7 (0.7)
ICD (or external cardioverter defibrillator)	24 (2.5)
Cardiac arrest location	
Home	603 (61.3)
Workplace	15 (1.5)
Public area	120 (12.2)
Nursing home	36 (3.7)
Ambulance	148 (15.1)
Hospital	40 (4.1)
Others	21 (2.1)
Witnessed arrest	776 (83.9)
Transport with EMS	975 (97.4)
Instructions from EMS dispatcher	98 (15.5)
Prehospital CPR	
None	105 (10.6)
Health-care provider	859 (86.5)
Bystander CPR	29 (2.9)
Prehospital CPR duration (min)	15 (8–20)
Time (min) from the arrest to the CPR	5 (1–10)
Initial cardiac arrest rhythm	
Asystole	682 (78.1)
PEA	97 (11.1)
VF	85 (9.7)
pVT	9 (1)
Shockable rhythm	
pVT/VF	94 (10.8)

Contd...

**Table 1: Contd...**

Variables	n (%) or mean±SD or median (25%–75%)
Asystole/PEA	779 (89.2)
Prehospital defibrillation	118 (12.1)
Number of prehospital defibrillations	2 (1–3)
Time (min) from the arrest to the first defibrillation	6.5 (3–12.75)
Prehospital airway procedure	
Not performed	117 (11.7)
BVM	318 (31.7)
ETI - successful	477 (47.6)
Supraglottic	54 (5.4)
Surgical	4 (0.4)
Unsuccessful	32 (3.2)
Prehospital chest compression	888 (89.4)
Prehospital ventilation support	899 (90.8)
Prehospital mechanical CPR device	9 (0.9)
Prehospital venous access	
None	112 (11.2)
Peripheral	886 (88.4)
Intraosseous	2 (0.2)
Central	2 (0.2)
Prehospital vascular access - exists	890 (88.8)
Prehospital epinephrine	741 (76.6)
Prehospital epinephrine dose (mg)	3 (2–5)
Prehospital amiodarone	25 (2.5)
Prehospital lidocaine	1 (0.1)
Prehospital sodium bicarbonate	5 (0.5)
Prehospital calcium	2 (0.2)
Reported by EMS before ED admission	710 (75.3)
Presence of pulse at ED admission - ROSC	136 (13.6)
Pulse duration in ED admission, if exists (min)	
<20	100 (77.5)
≥20	29 (22.5)
Survived event	
No	933 (93.1)
Yes	69 (6.9)

SD: Standard deviation, ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardiopulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, BVM: Bag-valve mask, ETI: Endotracheal intubation, ED: Emergency department, ROSC: Return of the spontaneous circulation

analyses between the two groups for the outcomes of survived event, overall sustained ROSC, survival to hospital discharge, and discharge with good neurological outcome are provided as supplemental files [Supplementary Tables 1-8]. It has been determined that the survival to hospital discharge rate is higher in male gender and patients with shockable rhythm and is lower, especially in the group of home localization and with mechanical chest compression devices [Supplementary Table 5]. In addition, it has been determined that both survived event, survival and good neurological outcome rates were higher in cases where bystander CPR was performed [Supplementary Tables 1, 5, and 7].

**Table 2: Descriptive variables of emergency department period**

Variables	n (%) or median (25%–75%)
First monitored rhythm at ED	
Asystole	682 (68.3)
PEA	124 (12.4)
VF	75 (7.5)
pVT	4 (0.4)
Pulse exists (any rhythm)	113 (11.3)
Shockable rhythm	
pVT/VF	79 (7.9)
Asystole/PEA	806 (80.8)
Pulse exists (any rhythm)	113 (11.3)
Advanced airway in ED	
Exists at admission	432 (43.1)
BVM	8 (0.8)
ETI	547 (54.6)
Supraglottic	7 (0.7)
Surgical	8 (0.8)
Venous access in ED	
None	1 (0.1)
Peripheral	943 (94.1)
Intraosseous	6 (0.6)
Central	52 (5.2)
Cardiac arrest again in ED	
No	232 (23.3)
Yes	218 (21.9)
No ROSC in ED	547 (54.9)
CPR time (min) to the first ROSC in ED	13 (7.5–20)
CPR in ED	950 (94.8)
Mechanical CPR device in ED	416 (41.5)
Defibrillation in ED	229 (22.9)
Number of defibrillations before first ROSC in ED	2 (1–3)
Total number of defibrillations in ED	3 (1–4)
Epinephrine in ED	941 (93.9)
Epinephrine dose before the first ROSC in ED (mg)	6 (3–10)
Total epinephrine dose in ED (mg)	6 (7–15)
Amiodarone in ED	157 (15.7)
Lidocaine in ED	5 (0.5)
Magnesium sulfate in ED	21 (2.1)
Sodium bicarbonate in ED	251 (25)
Calcium in ED	139 (13.9)
USG use during CPR in ED	522 (52.1)
Cardiac activity detected by USG	172 (33.1)
EtCO <sub>2</sub> device use during CPR in ED	302 (30.1)
5 <sup>th</sup> min ETCO <sub>2</sub> during CPR (mmHg)	14 (10–21.5)
Last ETCO <sub>2</sub> during CPR (mmHg)	14 (8–22)
Highest ETCO <sub>2</sub> during CPR (mmHg)	22 (15–34)
Initial pH in ED	6.98 (6.85–7.08)
Initial lactate (mmol/L) in ED	11.80 (8.04–15.40)
Outcome in ED	
Exitus	724 (72.3)
Ward/ICU	212 (21.2)
Transferred to other hospitals	66 (6.6)

Contd...

**Table 2: Contd...**

Variables	n (%) or median (25%–75%)
Outcome in hospital	
Exitus	958 (95.6)
Discharged	44 (4.4)
Neurological outcome at discharge	
Poor	967 (97.3)
Good	27 (2.7)
30-day survival	
Exitus	951 (94.9)
Survived	51 (5.1)
30-day neurological outcome	
Poor	966 (97.2)
Good	28 (2.8)
ST-segment elevation after ROSC	144 (34.4)
Coronary revascularization after ROSC	100 (23.9)
TTM after ROSC	9 (2.2)
Vasoactive drug after ROSC	271 (65.1)
ECLS after ROSC	1 (0.2)
IABP after ROSC	0
Surgery after ROSC	8 (2.0)
First pH after ROSC	7.05 (6.92–7.22)
First lactate (mmol/L) after ROSC	9.85 (5.59–13.93)
ROSC any (overall)	453 (45.2)
Sustained ROSC (overall)	334 (33.4)
Forensic case report	100 (10.0)

PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, BVM: Bag-valve mask, ETI: Endotracheal intubation, ROSC: Return of the spontaneous circulation, ED: Emergency department, med: median, CPR: Cardiopulmonary resuscitation, min: minute, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, ICU: Intensive care unit, TTM: Targeted temperature management, ECLS: Extracorporeal Life Support, IABP: Intra-aortic balloon pump

According to regional distribution, the highest bystander CPR rates were found in the Marmara and Aegean regions and the highest survival rates in the Mediterranean and Black Sea regions. The good neurological outcome rate is higher in the Mediterranean region [Supplementary Table 9].

### Multiple logistic regression analysis

As a result of multiple regression analyses, male gender (odds ratio [OR]: 2.691; 95% confidence interval [CI]: 1.186–6.103), prehospital initial shockable rhythm (OR: 6.480; 95% CI: 3.055–13.744), a shorter prehospital duration of CPR (OR: 0.919; 95% CI: 0.880–0.961), and the lack of CPR requirement in the ED (OR: 12.038; 95% CI: 5.000–28.984) were found to be positive independent predictors for the survival to hospital discharge [Table 3]. For sustained ROSC, presence of chronic pulmonary disease (OR: 1.677; 95% CI: 1.088–2.585), witnessed arrest (OR: 2.910; 95% CI: 1.588–5.336), arrest localization being hospital (OR: 3.703; 95% CI: 1.310–10.467; reference: home), shorter prehospital CPR duration (OR: 0.954; 95% CI: 0.933–0.975), not using mechanical chest compression device in ED (OR: 2.598; 95% CI: 1.774–3.803), the venous route used in the ED

**Table 3: Multiple regression analysis for survival to hospital discharge**

Variables	OR	95% CI	P
Day shift versus night shift and holidays	1.992	1.013–3.918	0.053
Gender (male vs. female)	2.691	1.186–6.103	0.018
Transport with EMS (EMS vs. not)	0.444	0.113–1.742	0.244
Prehospital initial shockable rhythm versus others	6.480	3.055–13.744	<0.001
Prehospital CPR duration (min)	0.919	0.880–0.961	<0.001
CPR in ED (none vs. performed)	12.038	5.000–28.984	<0.001
Sodium bicarbonate in ED (none vs. performed)	2.590	0.871–7.704	0.087

CI: Confidence interval, EMS: Emergency medical services, CPR: Cardiopulmonary resuscitation, ED: Emergency department, OR: Odds ratio

**Table 4: Multiple regression analysis for overall sustained return of the spontaneous circulation**

Variables	OR	95% CI	P
Chronic pulmonary disease	1.677	1.088–2.585	0.019
Witnessed arrest (yes vs. no)	2.910	1.588–5.336	0.001
Transport with EMS (EMS vs. non-EMS)	0.764	0.263–2.222	0.622
Cardiac arrest location (reference: Home)			
Workplace/public area	0.594	0.335–1.055	0.076
Nursing home	0.853	0.298–2.444	0.767
Ambulance	0.987	0.605–1.609	0.958
Hospital	3.703	1.310–10.467	0.014
Others	0.797	0.266–2.386	0.685
Prehospital CPR duration (min)	0.954	0.933–0.975	<0.001
Shockable rhythm at ED admission (reference: pVT/VF)			
Asystole/PEA	0.517	0.257–1.043	0.066
Pulse exists	2.031	0.840–4.911	0.116
Advanced airway in ED (ref: Exists at admission)			
BVM/supraglottic/surgical	0.151	0.017–1.304	0.086
ETI	0.824	0.555–1.225	0.339
Mechanical CPR device in ED (not performed vs. performed)	2.598	1.774–3.803	<0.001
Defibrillation in ED (performed vs. not)	1.583	0.984–2.547	0.058
Venous access in ED (others vs. peripheral)	2.105	1.051–4.219	0.036
Initial pH during CPR in ED (reference: pH<6.9)			
$6.9 \leq \text{pH} < 7.0$	2.106	1.257–3.528	0.005
$\text{pH} \geq 7.0$	3.005	1.941–4.652	<0.001

CI: Confidence interval, EMS: Emergency medical services, CPR: Cardiopulmonary resuscitation, ED: Emergency department, pVT: Pulseless ventricular tachycardia, VF: Ventricular fibrillation, PEA: Pulseless electrical activity, BVM: Bag-valve mask, ETI: Endotracheal intubation

being other than peripheral intravenous route (OR: 2.105; 95% CI: 1.051–4.219), and higher pH levels (for “ $6.9 \leq \text{pH} < 7.0$ ” OR: 2.106; 95% CI: 1.257–3.528 and for “ $\text{pH} \geq 7.0$ ” OR: 3.005; 95% CI: 1.941–4.652) were determined as independent predictors [Table 4].

## Discussion

This descriptive prospective multicenter study investigated the epidemiology and outcomes of OHCA patients in prehospital and emergency settings across Türkiye and aims to provide a comprehensive overview of this critical patient population. Of the OHCA patients, 453 (45.2%) achieved any ROSC, 334 (33.4%) achieved sustained ROSC, and 278 (27.7%) had survived event, which were the outcomes that primarily pertain to prehospital and emergency medicine department settings. Upon reviewing our findings, it can be concluded that our results closely align with

data reported in European and Australian studies. The EuRoCa TWO study, published in 2020, documented that any ROSC was achieved in 32.7% of 25,171 patients who experienced cardiac arrest and received CPR initiated by EMS or bystanders.<sup>[1]</sup> Similar findings have also been reported (23.8-37.8% for prehospital ROSC) in Australian and New Zealand data published by Beck *et al.*<sup>[8]</sup>

In a meta-analysis that examines 141 studies and nearly 4.6 million OHCA patients, the rate of sustained ROSC was reported as 29.7%. The authors conducted subgroup analyses by region and observed the highest sustained ROSC rate in Oceania countries (38.6%), followed by Europe (36.7%), and then Asian countries (22.1%).<sup>[9]</sup> While our sustained ROSC rates may not match those observed in Oceania and Europe, it can be inferred that our results are closely approximated. In the meta-analysis by Yan *et al.*, the global incidence of survived event rates among OHCA patients was reported as 22%. Oceanic countries exhibited the highest rate of survived events at 33.5%,





observed that better results were obtained in the male gender in terms of survival to hospital discharge and neurological outcome, but in multivariate analysis, the adjusted odds ratios were seen to be in favor of the female gender.<sup>[11]</sup> On the contrary, we observed better outcomes in male patients; we can attribute these results to the higher rates of bystander CPR in male gender.

The presence of an initial shockable rhythm as an independent predictor of survival to hospital discharge has been observed in various studies in different settings.<sup>[14-17]</sup> Additionally, in Li *et al.*'s data from Singapore, which shares similar characteristics and settings with our study and demonstrates a survival rate of 3.4%, shockable initial rhythm and shorter prehospital CPR duration were also identified as independent predictors.<sup>[14]</sup> It is evident that increasing the rate of bystander-initiated CPR can lead to improved outcome rates, highlighting the necessity for a nationwide educational campaign.

The presence of chronic pulmonary disease, witnessed arrest, cardiac arrest occurring at hospital perimeters, brief prehospital CPR duration, absence of mechanical CPR device uses in the ED, utilization of a venous access other than the peripheral intravenous route, and higher pH levels in the first blood gas obtained during CPR were identified as independent predictors of sustained ROSC.

A recent study by Balan *et al.* reported that witnessed cardiac arrest, initial shockable rhythm, and cardiac arrest occurring outside of home were found to be the independent predictors of hospital discharge in 3952 OHCA patients who achieved sustained ROSC.<sup>[16]</sup> Although our study populations and outcomes may differ, it is noteworthy that we obtained similar results.

The debate regarding whether mechanical or manual chest compressions yield different outcomes in cardiac arrest patients remains a hot topic of current research. Only a single 2015 randomized controlled study reported significant harm of mechanical compression device in OHCA patients in terms of survival with favorable neurological outcome.<sup>[18]</sup> A 2018 Cochrane review categorized existing studies as of medium-to-low quality and concluded that both methods were not superior to each other except in certain circumstances. It is important to note that this review included evaluations of both in-hospital and OHCA patients, with trauma patients excluded.<sup>[19]</sup> Therefore, our findings may not align precisely with those of the aforementioned review.

We found that the sustained ROSC rate was significantly higher in patients who received prehospital respiratory support, particularly among those who underwent successful ETI before arrival at the hospital. However,

the observational design of our study limits our ability to make definitive conclusions based solely on these findings. For instance, the recent AIRWAY-2 study investigated the impact of prehospital ETI and supraglottic airway (SGA) placement on 30-day survival in OHCA patients, finding no significant difference between the two methods.<sup>[20]</sup>

In our study, we observed a significantly higher rate of sustained ROSC among patients who received epinephrine prior to hospital arrival. This finding aligns with the PARAMEDIC2 study, a large-scale investigation conducted in 2018, which demonstrated higher rates of ROSC, survival to hospital admission, and 30-day survival in OHCA patients treated with prehospital epinephrine administration.<sup>[21]</sup> Consequently, the use of epinephrine in OHCA patients has gained stronger support, as reflected in the 2023 update of the AHA advanced cardiac life support guidelines.<sup>[22]</sup>

### Limitations

First, it should be noted that significant missing data exist, and consecutive sampling could not be properly implemented. Disruption of the operational functioning of study centers due to the earthquake on February 6, 2023, was a major contributing factor to these missing data. Additionally, two of the study centers had to be excluded from the study due to data unavailability. However, it has not been assessed whether this situation negatively impacts the regional representation of the country. The inclusion of tertiary hospitals in the study may introduce bias when extrapolating the results to the general public and rural ED.

Patients who received CPR outside and were not brought to the study centers were not included in the study; excluding these data also increases the bias factor. Most previous registry studies included only patients transported by EMS.<sup>[1,8-10,12]</sup> Although the rate of patients transported by non-EMS people is very low (2.6%), the inclusion of this patient group should be taken into consideration when discussing results with previous literature. Since reliable data could not be obtained about the time for prehospital transport, it was not included in the analysis. In most of the patients, the possible etiology of cardiac arrest was estimated by the physician based on the available history. Lack of knowledge about exact no-flow time (for lay person) may have a negative influence on results.

Another point, the hospital arrest localization group causes confusion for the results. In this group, cardiac arrest occurs at the triage or the hospital entrance. Therefore, it is understood that interventions such as prehospital chest compressions and respiratory support are not applied in a significant part of this group, and it

is understood that the prognosis is better, probably due to rapid access to health-care providers.

Given the study's nonrandomized controlled trial design, it is important to acknowledge that certain subgroup analysis findings may be influenced by confounding factors. For instance, while the lower success rate of CPR with mechanical devices might appear accurate, it is plausible that this outcome could be influenced by factors such as the preference for mechanical CPR devices in patients with a lower life expectancy and longer CPR duration.

## Conclusion

This study is the first multicenter, prospective study conducted in Türkiye revealing OHCA data. As a result, prognostic data obtained in the prehospital and ED periods are similar to those in developed countries; however, survival and good neurological outcome rates appear to be worse. Concepts such as ED crowding and intensive care quality can be discussed here. However, success in hospital discharge and neurological survival is an issue that requires multidisciplinary coordination. Although it is not the focus of this study, the correct approach would be to discuss these poor outcomes by focusing on the postcardiac arrest care step in the chain of survival.

Consistent with the literature, shockable initial rhythm and witnessed arrest were found to be associated with good outcomes. On the other hand, the bystander CPR rate, which is very low compared to developed countries, should be particularly scrutinized and efforts should be made to improve it by developing health policies and social policies.

### Acknowledgments

The researchers would like to thank the Emergency Medicine Association of Türkiye Resuscitation Study Group Members for their valuable contribution to the study.

### Author contributions

All authors conducted the study concept and design and acquisition of the data. AŞ and SK conducted the statistical analysis and interpretation of the data. AŞ and MMİ conducted the drafting of the manuscript. All authors conducted critical revision of the manuscript for important intellectual content.

### Conflicts of interest

None declared.

### Ethical approval

This multicenter study was carried out in 28 health centers in Türkiye within the scope of the Emergency Medicine Association of Türkiye (EMAT) Resuscitation Study Group, and approval was obtained from the Kocaeli University Non-Interventional Clinical Research Ethics Committee (KÜ GOKAEK-2022/18.24; November 10, 2022).

### Funding

None.

## References

- Gräsner JT, Wnent J, Herlitz J, Perkins GD, Lefering R, Tjelmeland I, et al. Survival after out-of-hospital cardiac arrest in Europe – Results of the EuReCa TWO study. *Resuscitation* 2020;148:218-26.
- Panchal AR, Bartos JA, Cabañas JG, Donnino MW, Drennan IR, Hirsch KG, et al. Part 3: Adult basic and advanced life support: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2020;142:S366-468.
- Merchant RM, Topjian AA, Panchal AR, Cheng A, Aziz K, Berg KM, et al. Part 1: Executive summary: 2020 American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2020;142:S337-57.
- Tekin FC, Köylü R, Köylü Ö, Kunt M. Factors related to resuscitation success and prognosis of cardiopulmonary arrest cases. *Indian J Crit Care Med* 2023;27:26-31.
- Halhalli HC, Şancı E, Uslu T. The comparison of manual and mechanical chest compression on survival and long-term neurological outcome of nontraumatic out-of-hospital cardiac arrest patients. *J Emerg Med* 2020;59:680-6.
- Şener A, Pamukçu Günaydın G, Tanrıverdi F. Mechanical versus manual chest compression: A retrospective-cohort in out-of-hospital cardiac arrest. *Acta Med* 2021;52:325-31.
- Perkins GD, Jacobs IG, Nadkarni VM, Berg RA, Bhanji F, Biarent D, et al. Cardiac arrest and cardiopulmonary resuscitation outcome reports: Update of the Utstein resuscitation registry templates for out-of-hospital cardiac arrest: A statement for healthcare professionals from a task force of the International Liaison Committee on Resuscitation (American Heart Association, European Resuscitation Council, Australian and New Zealand Council on Resuscitation, Heart and Stroke Foundation of Canada, Inter American Heart Foundation, Resuscitation Council of Southern Africa, Resuscitation Council of Asia); and the American Heart Association Emergency Cardiovascular Care Committee and the Council on Cardiopulmonary, Critical Care, Perioperative and Resuscitation. *Circulation* 2015;132:1286-300.
- Beck B, Bray J, Cameron P, Smith K, Walker T, Grantham H, et al. Regional variation in the characteristics, incidence and outcomes of out-of-hospital cardiac arrest in Australia and New Zealand: Results from the Aus-ROC epistry. *Resuscitation* 2018;126:49-57.
- Yan S, Gan Y, Jiang N, Wang R, Chen Y, Luo Z, et al. The global survival rate among adult out-of-hospital cardiac arrest patients who received cardiopulmonary resuscitation: A systematic review and meta-analysis. *Crit Care* 2020;24:61.
- Hawkes C, Booth S, Ji C, Brace-McDonnell SJ, Whittington A, Mapstone J, et al. Epidemiology and outcomes from out-of-hospital cardiac arrests in England. *Resuscitation* 2017;110:133-40.
- Kotini-Shah P, Del Rios M, Khosla S, Pugach O, Vellano K, McNally B, et al. Sex differences in outcomes for out-of-hospital cardiac arrest in the United States. *Resuscitation* 2021;163:6-13.
- Okubo M, Kiyohara K, Iwami T, Callaway CW, Kitamura T. Nationwide and regional trends in survival from out-of-hospital cardiac arrest in Japan: A 10-year cohort study from 2005 to 2014. *Resuscitation* 2017;115:120-8.
- Mentzelopoulos SD, Bossaert L, Raffay V, Askitopoulou H, Perkins GD, Greif R, et al. A survey of key opinion leaders on ethical resuscitation practices in 31 European countries. *Resuscitation* 2016;100:11-7.
- Li T, Yap J, Chng WQ, Tay JC, Shahidah N, Yeo C, et al. Clinicopathological correlates of out-of-hospital cardiac arrests. *J Arrhythm* 2022;38:416-24.
- Whittaker A, Lehal M, Calver AL, Corbett S, Deakin CD, Gray H, et al. Predictors of inhospital mortality following out-of-hospital cardiac arrest: Insights from a single-centre consecutive case

- series. *Postgrad Med J* 2016;92:250-4.
16. Balan P, Hsi B, Thangam M, Zhao Y, Monlezun D, Arain S, et al. The cardiac arrest survival score: A predictive algorithm for in-hospital mortality after out-of-hospital cardiac arrest. *Resuscitation* 2019;144:46-53.
  17. Baldi E, Caputo ML, Savastano S, Burkart R, Klersy C, Benvenuti C, et al. An utstein-based model score to predict survival to hospital admission: The UB-ROSC score. *Int J Cardiol* 2020;308:84-9.
  18. Perkins GD, Lall R, Quinn T, Deakin CD, Cooke MW, Horton J, et al. Mechanical versus manual chest compression for out-of-hospital cardiac arrest (PARAMEDIC): A pragmatic, cluster randomised controlled trial. *Lancet* 2015;385:947-55.
  19. Wang PL, Brooks SC. Mechanical versus manual chest compressions for cardiac arrest. *Cochrane Database Syst Rev* 2018;8:CD007260.
  20. Bengier JR, Kirby K, Black S, Brett SJ, Clout M, Lazaroo MJ, et al. Effect of a strategy of a supraglottic airway device versus tracheal intubation during out-of-hospital cardiac arrest on functional outcome: The AIRWAYS-2 randomized clinical trial. *JAMA* 2018;320:779-91.
  21. Perkins GD, Ji C, Deakin CD, Quinn T, Nolan JP, Scomparin C, et al. A randomized trial of epinephrine in out-of-hospital cardiac arrest. *N Engl J Med* 2018;379:711-21.
  22. Perman SM, Elmer J, Maciel CB, Uzendu A, May T, Mumma BE, et al. 2023 American Heart Association Focused Update on Adult Advanced Cardiovascular Life Support: An update to the American Heart Association guidelines for cardiopulmonary resuscitation and emergency cardiovascular care. *Circulation* 2024;149:e254-73.

**Supplementary Table 1: Factors associated with survived event**

Variables	Survived event, n (%)	P
Time zone		
Day shift	24 (6.2)	0.594
Night shift and holidays	45 (7.3)	
Gender		
Female	34 (8.7)	0.068
Male	35 (5.7)	
Independent living		
No	13 (6.2)	0.709
Yes	52 (7.2)	
Chronic neurological disease		
No	56 (7.3)	0.666
Yes	10 (6.0)	
Chronic pulmonary disease		
No	57 (7.5)	0.425
Yes	10 (5.5)	
Chronic cardiovascular disease		
No	26 (5.8)	0.242
Yes	38 (7.7)	
Active malignancy		
No	61 (7.5)	0.212
Yes	5 (4.0)	
Ventricular assist device		
No	66 (6.9)	0.399
Yes	1 (14.3)	
ICD (or external cardioverter defibrillator)		
No	65 (6.9)	0.680
Yes	2 (8.3)	
Witnessed arrest		
Yes	59 (7.6)	0.429
No	8 (5.4)	
Instructions from EMS dispatcher		
Yes	8 (8.2)	0.880
No	38 (7.1)	
Prehospital CPR		
None	1 (1)	0.016
Health-care provider	63 (7.3)	
Bystander CPR	4 (13.8)	
Initial arrest rhythm		
Asystole	46 (6.7)	0.811
PEA	5 (5.2)	
VF	7 (8.2)	
pVT	1 (11.1)	
Shockable rhythm		
pVT/VF	8 (8.5)	0.618
Asystole/PEA	51 (6.5)	
Prehospital defibrillation		
No	58 (6.8)	0.408
Yes	11 (9.3)	
Prehospital chest compression		
None	1 (1)	0.020
Performed	67 (7.5)	

**Supplementary Table 1: Contd...**

Variables	Survived event, n (%)	P
Prehospital ventilation support		
None	2 (2.2)	0.097
Performed	67 (7.5)	
Prehospital mechanical CPR device		
None	66 (6.7)	0.122
Used	2 (22.2)	
Prehospital vascular access		
None	2 (1.8)	0.039
Yes	67 (7.5)	
Prehospital epinephrine		
None	4 (1.8)	0.001
Performed	63 (8.5)	
Prehospital amiodarone		
None	65 (6.8)	0.685
Performed	2 (8)	
Prehospital lidocaine		
None	67 (6.8)	0.069
Performed	1 (100)	
Prehospital sodium bicarbonate		
None	66 (6.7)	0.041
Performed	2 (40)	
Prehospital calcium		
None	68 (6.9)	1.000
Performed	-	
Reported by EMS before ED admission		
Yes	47 (6.6)	1.000
No	15 (6.4)	
Etiology		
Acute coronary syndrome	30 (6.5)	0.129
Pulmonary embolism	6 (14)	
Electrolyte disorder	3 (5.9)	
Asthma	3 (18.8)	
Drowning	3 (18.8)	
Others, unknown	24 (5.8)	
Prehospital airway procedure		
Not performed	3 (2.6)	<0.001*
BVM	13 (4.1)	
ETI - successful/surgical	51 (10.6)	
Supraglottic	1 (1.9)	
Unsuccessful	1 (3.1)	
Cardiac arrest location		
Home	43 (7.1)	0.239
Workplace/public area	13 (9.6)	
Nursing home	3 (8.3)	
Ambulance	6 (4.1)	
Hospital	3 (7.5)	
Others	-	

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation

Contd...

**Supplementary Table 2: Factors associated with survived event - continues variables**

<b>Variables</b>	<b>Nonsurvived event</b>	<b>Survived event</b>	<b>P*</b>
Age (year)	67.1±15.1	65.9±15.7	0.655
Prehospital CPR duration (min)	15 (8–20)	15 (5.3–20)	0.897
Time from the arrest to the CPR (min)	5 (1–10)	10 (2–10)	0.714
Number of defibrillation (if exist)	2 (1–3)	2 (1–3)	0.712
Time from the arrest to the first defibrillation (min)	6.5 (3–11.8)	6.5 (2.3–15.3)	0.966

\*Age: Independent samples *t*-test, mean±SD. Variables other than age: Mann–Whitney *U*-test, median (25%–75%). CPR: Cardio-pulmonary resuscitation, SD: Standard deviation

**Supplementary Table 3: Factors associated with sustained return of the spontaneous circulation (overall)**

Variables	Sustained ROSC, n (%)	P
Time zone		
Day shift	115 (29.9)	0.065
Night shift and holidays	219 (35.6)	
Gender		
Female	136 (35.1)	0.388
Male	198 (32.4)	
Independent living		
No	62 (29.5)	0.070
Yes	261 (36.3)	
Chronic neurological disease		
No	264 (34.3)	0.693
Yes	55 (32.7)	
Chronic pulmonary disease		
No	244 (32.1)	0.009
Yes	77 (42.3)	
Chronic cardiovascular disease		
No	154 (34.1)	0.708
Yes	163 (33.0)	
Active malignancy		
No	278 (34.1)	0.685
Yes	40 (32.3)	
Ventricular assist device		
No	320 (33.7)	0.049
Yes	5 (71.4)	
ICD (or external cardioverter defibrillator)		
No	320 (34.0)	1.000
Yes	8 (33.3)	
Witnessed arrest		
Yes	296 (38.2)	<0.001
No	26 (17.4)	
Instructions from EMS dispatcher		
Yes	32 (32.7)	0.625
No	187 (35.2)	
Prehospital CPR		
None	55 (52.4)	<0.001
Health-care provider	264 (30.8)	
Bystander CPR	11 (37.9)	
Prehospital initial cardiac arrest rhythm		
Asystole	206 (30.2)	<0.001
PEA	38 (39.6)	
VF	46 (54.1)	
pVT	3 (33.3)	
Prehospital shockable rhythm		
pVT/VF	49 (52.1)	<0.001
Asystole/PEA	244 (31.4)	
Prehospital defibrillation		
No	283 (33.1)	0.367
Yes	44 (37.3)	
Prehospital chest compression		
None	55 (52.4)	<0.001
Performed	275 (31.1)	
Prehospital ventilation support		
None	50 (54.9)	<0.001
Performed	282 (31.5)	

**Supplementary Table 3: Contd...**

Variables	Sustained ROSC, n (%)	P
Prehospital mechanical CPR device		
None	328 (33.4)	1.000
Used	3 (33.3)	
Prehospital vascular access		
None	47 (42.0)	0.042
Yes	287 (32.4)	
Prehospital epinephrine		
None	104 (46.2)	<0.001
Performed	226 (30.6)	
Prehospital amiodarone		
None	320 (33.5)	0.640
Performed	10 (40.0)	
Prehospital lidocaine		
None	330 (33.6)	0.337
Performed	1 (100)	
Prehospital sodium bicarbonate		
None	329 (33.6)	1.000
Performed	2 (40)	
Prehospital calcium		
None	331 (33.7)	0.553
Performed	-	
Reported by EMS before ED admission		
Yes	206 (29.1)	<0.001
No	112 (48.3)	
Etiology		
Drowning	11 (68.8)	<0.001
Acute coronary syndrome	156 (33.6)	
Pulmonary embolism	19 (46.3)	
Electrolyte disorder	16 (31.4)	
Asthma	12 (75)	
Others, unknown	120 (29.2)	
Prehospital airway procedure		
Not performed	64 (54.7)	<0.001
BVM	82 (25.9)	
ETI - successful/surgical	170 (35.5)	
Supraglottic	13 (24.1)	
Unsuccessful	5 (15.6)	
Cardiac arrest location		
Home	177 (29.5)	<0.001
Workplace/public area	42 (31.1)	
Nursing home	8 (22.2)	
Ambulance	66 (44.9)	
Hospital	31 (77.5)	
Others	8 (38.1)	
Shockable rhythm at ED admission		
pVT/VF	44 (55.7)	<0.001
Asystole/PEA	203 (25.3)	
Pulse exists	85 (75.2)	
Advanced airway in ED		
Exists at admission	146 (34)	0.111
BVM/Supraglottic/surgical	3 (13)	
ETI	185 (33.9)	
CPR in ED		
None	52 (100)	<0.001
Performed	282 (29.8)	

Contd...

Contd...

**Supplementary Table 3: Contd...**

<b>Variables</b>	<b>Sustained ROSC, n (%)</b>	<b>P</b>
Mechanical CPR device in ED		
None	236 (40.4)	<0.001
Used	98 (23.6)	
Defibrillation in ED		
None	234 (30.4)	<0.001
Performed	100 (43.7)	
Venous access in ED		
Others	32 (54.2)	<0.001
Peripheral	302 (32.1)	
Epinephrine in ED		
None	61 (100)	<0.001
Performed	273 (29.1)	
Amiodarone in ED		
None	263 (31.2)	0.001
Performed	71 (45.2)	
Lidocaine in ED		
None	332 (33.4)	1.000
Performed	2 (40)	
Magnesium sulfate in ED		
None	324 (33.1)	0.246
Performed	10 (47.6)	
Sodium bicarbonate in ED		
None	236 (31.5)	0.022
Performed	98 (39.4)	
Calcium in ED		
None	290 (33.7)	0.678
Performed	44 (31.9)	
USG use during CPR in ED		
None	156 (32.6)	0.578
Used	178 (34.2)	
If USG used, cardiac activity exists		
None	34 (9.8)	<0.001
Exists	143 (83.1)	
EtCO2 device use during CPR in ED		
None	257 (36.9)	<0.001
Used	77 (25.5)	

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation, USG: Ultrasonography, EtCO2: End-tidal carbon dioxide, ROSC: Return of the spontaneous circulation



**Supplementary Table 4: Factors associated with sustained return of the spontaneous circulation (overall) - continues variables**

Variables	No sustained ROSC	Sustained ROSC	<i>P</i> *
Age (year)	67.4±15.4	66.2±14.9	0.242
Prehospital CPR duration (min)	15 (10–25)	10 (3–20)	<0.001
Prehospital time from the arrest to the CPR (min)	8 (2–10)	5 (1–10)	<0.001
Prehospital number of defibrillation (if exist)	2 (1–3)	2 (1–3)	0.967
Prehospital time from the arrest to the first defibrillation (min)	10 (5.5–15)	4 (1–9)	0.001
Number of defibrillations before first ROSC in ED	2 (1–3)	3 (1–4)	0.074
Total number of defibrillations in ED	3 (1–4)	3 (1–4)	0.648
Epinephrine dose (mg) before the first ROSC in ED	10 (5–14)	4 (2–6)	<0.001
Total epinephrine dose (mg) in ED	13 (10–15)	6 (3–11)	<0.001
CPR to the first ROSC in ED (min)	12.5 (5.5–21.8)	14 (8–20)	0.843
Initial pH during CPR in ED	6.93 (6.80–7.05)	7.02 (6.92–7.13)	<0.001
Initial lactate during CPR in ED (mmol/L)	12.5 (9–16)	10.24 (6.8–13.2)	<0.001
5 <sup>th</sup> min ETCO <sub>2</sub> during CPR in ED (mmHg)	12 (8.5–18)	18 (12–29.3)	<0.001
Last ETCO <sub>2</sub> during CPR in ED (mmHg)	11 (6.75–16)	35 (24–54)	<0.001
Highest ETCO <sub>2</sub> during CPR in ED (mmHg)	18 (13–25)	39 (30.5–55)	<0.001

\*Age: Independent samples *t*-test, mean±SD. Variables other than age: Mann–Whitney *U*-test; median (25%–75%). ROSC: Return of the spontaneous circulation, CPR: Cardio-pulmonary resuscitation, ED: Emergency department, ETCO<sub>2</sub>: End-tidal carbon dioxide, SD: Standard deviation

**Supplementary Table 5: Factors associated with survival to hospital discharge**

Variables	Survival to hospital discharge, n (%)	P
Time zone		
Day shift	25 (6.5)	0.017
Night shift and holidays	19 (3.1)	
Gender		
Female	2 (2.3)	0.016
Male	35 (5.7)	
Independent living		
No	4 (1.9)	0.044
Yes	40 (5.5)	
Chronic neurological disease		
No	37 (4.8)	0.629
Yes	6 (3.6)	
Chronic pulmonary disease		
No	37 (4.9)	0.469
Yes	6 (3.3)	
Chronic cardiovascular disease		
No	27 (6)	0.061
Yes	16 (3.2)	
Active malignancy		
No	42 (5.1)	0.051
Yes	1 (0.8)	
Ventricular assist device		
No	44 (4.6)	1.000
Yes	-	
ICD (or external cardioverter defibrillator)		
No	44 (4.7)	0.622
Yes	-	
Witnessed arrest		
Yes	37 (4.8)	0.587
No	5 (3.4)	
Instructions from EMS dispatcher		
Yes	2 (2)	0.406
No	24 (4.5)	
Prehospital CPR		
None	11 (10.5)	0.001
Health-care provider	29 (3.4)	
Bystander CPR	4 (13.8)	
Prehospital shockable rhythm		
pVT/VF	17 (18.1)	<0.001
Asystole/PEA	19 (2.4)	
Prehospital defibrillation		
No	31 (3.6)	0.003
Yes	12 (10.2)	
Prehospital chest compression		
None	11 (10.5)	0.004
Performed	33 (3.7)	
Prehospital ventilation support		
None	12 (13.2)	<0.001
Performed	32 (3.6)	
Prehospital mechanical CPR device		
None	43 (4.4)	1.000
Used	-	

**Supplementary Table 5: Contd...**

Variables	Survival to hospital discharge, n (%)	P
Prehospital vascular access		
None	9 (8)	0.081
Yes	35 (3.9)	
Prehospital epinephrine		
None	21 (9.3)	<0.001
Performed	23 (3.1)	
Prehospital amiodarone		
None	43 (4.5)	1.000
Performed	1 (4)	
Prehospital lidocaine		
None	44 (4.5)	1.000
Performed	-	
Prehospital sodium bicarbonate		
None	44 (4.5)	1.000
Performed	-	
Prehospital calcium		
None	44 (4.5)	1.000
Performed	-	
Survival event		
None	32 (3.4)	<0.001
Yes	12 (17.4)	
Reported by EMS before ED admission		
Yes	25 (3.5)	0.006
No	19 (8.2)	
Etiology		
Drowning	2 (12.5)	0.085
Acute coronary syndrome	28 (6)	
Pulmonary embolism	1 (2.3)	
Electrolyte disorder	1 (2)	
Asthma	-	
Others, unknown	12 (2.9)	
Prehospital airway procedure		
Not performed	17 (14.5)	<0.001
BVM	10 (3.1)	
ETI - successful/surgical	15 (3.1)	
Supraglottic	1 (1.9)	
Unsuccessful	1 (3.1)	
Cardiac arrest location		
Home	16 (2.7)	0.002
Workplace/public area	12 (8.9)	
Nursing home	-	
Ambulance	9 (6.1)	
Hospital	5 (12.5)	
Others	1 (4.8)	
Shockable rhythm at ED admission		
pVT/VF	13 (6.5)	<0.001
Asystole/PEA	17 (2.1)	
Pulse exists	14 (12.4)	
Advanced airway in ED		
Exists at admission	14 (3.2)	0.295
BVM/Supraglottic/Surgical	1 (4.3)	
ETI	29 (5.3)	

Contd...

Contd...

Supplementary Table 5: Contd...

Variables	Survival to hospital discharge, n (%)	P
CPR in ED		
None	12 (23.1)	<0.001
Performed	32 (3.4)	
Mechanical CPR device in ED		
None	33 (5.6)	0.034
Used	11 (2.6)	
Defibrillation in ED		
None	27 (3.5)	0.018
Performed	17 (7.4)	
Venous access in ED		
Others	3 (5.1)	0.740
Peripheral	41 (4.3)	
Epinephrine in ED		
None	14 (23)	<0.001
Performed	30 (3.2)	
Amiodarone in ED		
None	30 (3.6)	0.005
Performed	14 (8.9)	
Lidocaine in ED		
None	43 (4.3)	0.201
Performed	1 (20)	
Magnesium sulfate in ED		
None	40 (4.1)	0.011
Performed	4 (19)	
Sodium bicarbonate in ED		
None	40 (5.3)	0.020
Performed	4 (1.6)	
Calcium in ED		
None	43 (5)	0.040
Performed	1 (0.7)	
Cardiac arrest again in ED		
None	40 (17.2)	<0.001
Yes	4 (1.8)	
No ROSC in ED	-	
USG use during CPR in ED		
None	21 (4.4)	1.000
Used	23 (4.4)	
If USG used, cardiac activity exists		
None	2 (0.6)	<0.001
Exists	21 (12.2)	
EtCO <sub>2</sub> device use during CPR in ED		
None	34 (4.9)	0.353
Used	10 (3.3)	
ST segment elevation after ROSC		
No	21 (7.7)	0.014
Yes	23 (16)	
Coronary revascularization after ROSC		
No	17 (5.3)	<0.001
Yes	27 (27)	
TTM after ROSC		
No	39 (9.8)	0.055
Yes	3 (33.3)	

Supplementary Table 5: Contd...

Variables	Survival to hospital discharge, n (%)	P
Vasoactive drug after ROSC		
None	22 (15.2)	0.019
Performed	20 (7.4)	
ECLS after ROSC		
None	44 (10.7)	1.000
Performed	-	
Surgery after ROSC		
No	43 (10.7)	0.601
Yes	1 (12.5)	
Forensic case report		
None	39 (4.3)	0.795
Performed	5 (5)	

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation, ROSC: Return of the spontaneous circulation, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, TTM: Targeted temperature management, ECLS: Extracorporeal life support

Contd...

**Supplementary Table 6: Factors associated with survival to hospital discharge - continues variables**

Variables	Exitus	Discharged	P*
Age (year)	67.3±15.2	61.0±14.7	0.007
Prehospital CPR duration (min)	15 (10–20)	8.5 (0.13–12)	<0.001
Prehospital time from the arrest to the CPR (min)	5 (1–10)	3 (1–7.3)	0.036
Prehospital number of defibrillations (if exist)	2 (1–3)	1.5 (1–2)	0.353
Prehospital time from the arrest to the first defibrillation (min)	9 (4–15)	2 (0.8–4.3)	0.001
Number of defibrillations before first ROSC in ED	2 (1–3)	3 (1–5.5)	0.198
Total number of defibrillations in ED	3 (1–4)	3 (1–4.5)	0.711
Epinephrine dose (mg) before the first ROSC in ED	6.5 (3–10)	3 (2–5)	<0.001
Total epinephrine dose (mg) in ED	11 (8–15)	3 (2–4.8)	<0.001
CPR to the first ROSC in ED (min)	14.5 (8–22)	10 (5–12)	0.002
Initial pH during CPR in ED	6.97 (6.84–7.07)	7.11 (7–7.24)	<0.001
Initial lactate during CPR in ED (mmol/L)	12 (8.33–15.46)	7.98 (5.63–11.93)	<0.001
5 <sup>th</sup> min ETCO <sub>2</sub> during CPR in ED (mmHg)	14 (10–21)	16 (11–43.8)	0.148
Last ETCO <sub>2</sub> during CPR in ED (mmHg)	14 (8–20)	47.5 (33–55.8)	<0.001
Highest ETCO <sub>2</sub> during CPR in ED (mmHg)	20 (15–33.3)	49 (34.8–55.8)	<0.001
pH after ROSC in ED	7.03 (6.91–7.18)	7.25 (7.09–7.31)	<0.001
Lactate after ROSC in ED (mmol/L)	10.63 (6.33–14.13)	4.59 (3.08–7.91)	<0.001

\*Age: Independent Samples *t*-test; mean±SD. Variables other than age: Mann–Whitney *U*-test; median (25%–75%). CPR: Cardio-pulmonary resuscitation, ROSC: Return of the spontaneous circulation, ED: Emergency department, EtCO<sub>2</sub>: End-tidal carbon dioxide, SD: Standard deviation

**Supplementary Table 7: Factors associated with good neurological outcome at discharge**

Variables	Good neurological outcome at discharge, n (%)	P
Time zone		
Day shift	16 (4.2)	0.041
Night shift and holidays	11 (1.8)	
Gender		
Female	5 (1.3)	0.044
Male	22 (3.6)	
Independent living		
No	-	0.009
Yes	27 (3.8)	
Chronic neurological disease		
No	25 (3.3)	0.067
Yes	1 (0.6)	
Chronic pulmonary disease		
No	25 (3.3)	0.076
Yes	1 (0.6)	
Chronic cardiovascular disease		
No	17 (3.8)	0.100
Yes	9 (1.8)	
Active malignancy		
No	26 (3.2)	0.038
Yes	-	
Ventricular assist device		
No	27 (2.9)	1.000
Yes	-	
ICD (or external cardioverter defibrillator)		
No	27 (2.9)	1.000
Yes	-	
Witnessed arrest		
Yes	24 (3.1)	0.160
No	1 (0.7)	
Instructions from EMS dispatcher		
Yes	1 (1)	0.488
No	15 (2.8)	
Prehospital CPR		
None	9 (8.7)	<0.001
Health-care provider	14 (1.6)	
Bystander CPR	4 (13.8)	
Prehospital shockable rhythm		
pVT/VF	15 (16)	<0.001
Asystole/NEA	5 (0.6)	
Prehospital defibrillation		
No	16 (1.9)	<0.001
Yes	10 (8.5)	
Prehospital chest compression		
None	9 (8.7)	0.001
Performed	18 (2)	
Prehospital ventilation support		
None	9 (10)	<0.001
Performed	18 (2)	
Prehospital mechanical CPR device		
None	26 (2.7)	1.000
Used	-	

Contd...

**Supplementary Table 7: Contd...**

Variables	Good neurological outcome at discharge, n (%)	P
Prehospital vascular access		
None	6 (5.4)	0.110
Yes	21 (2.4)	
Prehospital epinephrine		
None	18 (8)	<0.001
Performed	9 (1.2)	
Prehospital amiodarone		
None	26 (2.7)	0.508
Performed	1 (4)	
Prehospital lidocaine		
None	27 (2.8)	1.000
Performed	-	
Prehospital sodium bicarbonate		
None	27 (2.8)	1.000
Performed	-	
Prehospital calcium		
None	27 (2.8)	1.000
Performed	-	
Survival event		
None	20 (2.2)	0.001
Yes	7 (10.4)	
Reported by EMS before ED admission		
Yes	14 (2)	0.008
No	13 (5.6)	
Etiology		
Drowning	2 (12.5)	<0.001
Acute coronary syndrome	22 (4.8)	
Pulmonary embolism	-	
Electrolyte disorder	1 (2)	
Asthma	-	
Others, unknown	2 (0.5)	
Prehospital airway procedure		
Not performed	14 (12.1)	<0.001
BVM	6 (1.9)	
ETI - successful/surgical	7 (1.5)	
Supraglottic	-	
Unsuccessful	-	
Cardiac arrest location		
Home	6 (1.0)	<0.001
Workplace/public area	8 (6)	
Nursing home	-	
Ambulance	6 (4.1)	
Hospital	5 (12.5)	
Others	1 (4.8)	
Shockable rhythm at ED admission		
pVT/VF	12 (15.2)	<0.001
Asystole/PEA	6 (0.8)	
Pulse exists	9 (8.1)	
Advanced airway in ED		
Exists at admission	7 (1.6)	0.188
BVM/supraglottic/surgical	1 (4.3)	
ETI	19 (3.5)	
CPR in ED		
None	7 (14)	<0.001
Performed	20 (2.1)	

Contd...

Supplementary Table 7: Contd...

Variables	Good neurological outcome at discharge, n (%)	P
Mechanical CPR device in ED		
None	20 (3.5)	0.133
Used	7 (1.7)	
Defibrillation in ED		
None	11 (1.4)	<0.001
Performed	16 (7)	
Venous access in ED		
Others	1 (1.7)	1.000
Peripheral	26 (2.8)	
Epinephrine in ED		
None	9 (15.3)	<0.001
Performed	18 (1.9)	
Amiodarone in ED		
None	15 (1.8)	<0.001
Performed	12 (7.7)	
Lidocaine in ED		
None	26 (2.6)	0.129
Performed	1 (20)	
Magnesium sulfate in ED		
None	23 (2.4)	0.002
Performed	4 (19)	
Sodium bicarbonate in ED		
None	25 (3.4)	0.054
Performed	2 (0.8)	
Calcium in ED		
None	26 (3)	0.160
Performed	1 (0.7)	
Cardiac arrest again in ED		
None	24 (10.7)	<0.001
Yes	3 (1.4)	
No ROSC in ED	-	
USG use during CPR in ED		
None	14 (3)	0.807
Used	13 (2.5)	
If USG used, cardiac activity exists		
None	-	<0.001
Exists	13 (7.6)	
EtCO2 device use during CPR in ED		
None	21 (3)	0.470
Used	6 (2)	
ST elevation after ROSC		
No	8 (3)	<0.001
Yes	19 (13.3)	
Coronary revascularization after ROSC		
No	5 (1.6)	<0.001
Yes	22 (22.2)	
TTM after ROSC		
No	24 (6.2)	0.445
Yes	1 (11.1)	
Vasoactive drug after ROSC		
None	17 (12.1)	0.001
Performed	9 (3.4)	

Contd...

Supplementary Table 7: Contd...

Variables	Good neurological outcome at discharge, n (%)	P
ECLS after ROSC		
None	27 (6.7)	1.000
Performed	-	
Surgery after ROSC		
No	26 (6.6)	0.430
Yes	1 (12.5)	
Forensic case report		
None	24 (2.7)	0.747
Performed	3 (3)	

Pearson's Chi-square test, Fisher's exact test. ICD: Implantable cardioverter-defibrillator, EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, PEA: Pulseless electrical activity, VF: Ventricular fibrillation, pVT: Pulseless ventricular tachycardia, ED: Emergency department, BVM: Bag-valve mask, ETI: Endotracheal intubation, ROSC: Return of the spontaneous circulation, USG: Ultrasonography, EtCO2: End-tidal carbon dioxide, TTM: Targeted temperature management, ECLS: Extracorporeal life support

**Supplementary Table 8: Factors associated with good neurological outcome at discharge - continuous variables**

Variables	Poor	Good	P*
Age (year)	67.3±15.2	57.3±12.4	0.001
Prehospital CPR duration (min)	15 (10–20)	2 (0–10)	<0.001
Prehospital time from the arrest to the CPR (min)	5 (1–10)	2 (0.5–5.5)	0.047
Prehospital number of defibrillations (if exist)	2 (1–3)	1.5 (1–2.5)	0.457
Prehospital time from the arrest to the first defibrillation (min)	9 (4–15)	2 (0.5–4.5)	0.002
Number of defibrillations before first ROSC in ED	2 (1–3)	3 (1–5.5)	0.198
Total number of defibrillations in ED	3 (1–4)	3 (1–4.8)	0.856
Epinephrine dose (mg) before the first ROSC in ED	6 (3–10)	3 (1.5–4.5)	<0.001
Total epinephrine dose (mg) in ED	11 (8–15)	4 (1.8–5)	<0.001
CPR to the first ROSC in ED (min)	14 (8–21.8)	8 (5–14.3)	0.009
Initial pH during CPR in ED	6.97 (6.84–7.07)	7.16 (7.04–7.27)	<0.001
Initial lactate during CPR in ED (mmol/L)	12 (8.33–15.42)	7.10 (5.30–11.80)	<0.001
5 <sup>th</sup> min ETCO <sub>2</sub> during CPR in ED (mmHg)	14 (10–21)	25.5 (12.5–47)	0.109
Last ETCO <sub>2</sub> during CPR in ED (mmHg)	14 (8–20)	43 (30–60.5)	0.001
Highest ETCO <sub>2</sub> during CPR in ED (mmHg)	21 (15–34)	44.5 (34.8–60.5)	0.002
pH after ROSC in ED	7.05 (6.91–7.19)	7.28 (7.09–7.33)	<0.001
Lactate after ROSC in ED (mmol/L)	10.35 (6.25–14)	4.05 (2.13–10.02)	<0.001

\*Age: Independent samples *t*-test, mean±SD. Variables other than age: Mann–Whitney *U*-test, median (25%–75%). CPR: Cardio-pulmonary resuscitation, ROSC: Return of the spontaneous circulation, ED: Emergency department, ETCO<sub>2</sub>: End-tidal carbon dioxide, SD: Standard deviation

**Supplementary Table 9: Distribution of outcomes in terms of regions**

	Regions						
	Marmara, n (%)	Aegean, n (%)	Mediterranean, n (%)	Central Anatolia, n (%)	Black Sea, n (%)	Eastern Anatolia, n (%)	Southeastern Anatolia, n (%)
EMS	378 (97.7)	326 (96.4)	101 (98.1)	62 (96.9)	39 (97.5)	40 (100.0)	29 (100.0)
Bystander CPR	14 (3.6)	12 (3.6)	2 (2.0)	0	0	1 (2.5)	0
Shockable rhythm	37 (11.5)	3 (11.5)	23 (24.2)	24 (8)	2 (5.1)	3 (8.1)	2 (3.8)
Prehospital airway procedure							
Not performed	39 (10.1)	2 (6.9)	20 (19.4)	40 (11.8)	4 (10)	2 (5.0)	10 (15.6)
BVM	134 (34.6)	3 (10.3)	31 (30.1)	93 (27.4)	11 (27.5)	25 (62.5)	21 (32.8)
ETI - successful	185 (47.8)	22 (75.9)	48 (46.6)	159 (46.9)	22 (55)	11 (27.5)	30 (46.9)
Supraglottic	12 (3.1)	2 (6.9)	3 (2.9)	33 (9.7)	0	1 (2.5)	3 (4.7)
Surgical	1 (0.3)	0	0	1 (0.3)	2 (5.0)	0	0
Unsuccessful	16 (4.1)	0	1 (1)	13 (3.8)	1 (2.5)	1 (2.5)	0
Prehospital venous access							
None	35 (9.0)	1 (3.4)	9 (8.7)	45 (13.3)	5 (12.5)	9 (22.5)	8 (12.5)
Peripheral	351 (90.7)	27 (93.1)	93 (90.3)	293 (86.4)	35 (87.5)	31 (77.5)	56 (87.5)
Intraosseous	1 (0.3)	1 (3.4)	0	0	0	0	0
Central	0	0	1 (1.0)	1 (0.3)	0	0	0
Mechanical CPR device in ED	180 (46.5)	8 (27.6)	10 (9.7)	170 (50.1)	26 (65.0)	19 (47.5)	3 (4.7)
USG use during CPR in ED	129 (33.3)	19 (65.5)	75 (72.8)	229 (67.6)	22 (55.0)	5 (12.5)	43 (67.2)
EtCO <sub>2</sub> device use during CPR in ED	96 (24.8)	6 (20.7)	46 (44.7)	136 (40.1)	16 (40.0)	2 (5.0)	0
Survived event	15 (3.9)	19 (5.6)	17 (16.5)	5 (7.8)	2 (5.0)	3 (7.5)	8 (27.6)
Sustained ROSC	102 (26.4)	114 (33.7)	59 (57.8)	21 (32.8)	12 (30.0)	12 (30.0)	14 (48.3)
Survival to hospital discharge	13 (3.4)	10 (2.9)	14 (13.6)	2 (3.1)	3 (7.5)	1 (2.5)	1 (3.4)
Good neurological outcome at discharge	11 (2.9)	4 (1.2)	11 (10.7)	0	0	1 (2.5)	0

EMS: Emergency medical service, CPR: Cardio-pulmonary resuscitation, BVM: Bag-valve mask, ETI: Endotracheal intubation, ED: Emergency department, USG: Ultrasonography, EtCO<sub>2</sub>: End-tidal carbon dioxide, ROSC: Return of the spontaneous circulation